



## **ADAPTATION MECHANISMS OF SOYBEAN TO DROUGHT AND ENVIRONMENTAL STRESS FACTORS**

**Fozilov Sherzod Musurmonovich**

Termez State University, Teacher. Surkhandarya, Uzbekistan

**Sulliyeva Suluv Xurramovna**

Termez State University, Associate Professor, Surkhandarya, Uzbekistan

**Abstract:** The growth and development of soybean and the quality and composition of the crop are highly influenced by external environmental factors. Each influencing factor differs from one another according to its characteristics. It should be said that the soybean plant does not choose a place according to its biological characteristics. It also grows in saline, cultivated fields with elevated groundwater levels. It is also very convenient to care for it in a mixed way.

**Key words:** genotypic, mesophyte, critical phase, nodule, ripening phase, photosynthesis, transpiration.

Depending on the conditions of origin of the soybean plant, with genotypic resistance to drought and high humidity, cold and high temperatures, infertile and high levels of mineral fertilizers, dense and porous soils, as well as salty and sour. resistant to soils, but when these factors are optimized, it grows well, develops and gives high yields.

*Drought resistance* . Among legumes, soybean is second only to pea in terms of drought tolerance. Drought resistance is related to the centers of origin, Korean and Indian subspecies are less drought resistant than Manchurian and Chinese species. In the first half of the shadow period, the above-ground part grows slowly, and during the period of active



growth of the root system, it is resistant to drought (until the flowering period). Water requirements are very demanding during the flowering, podding and seeding periods. When there is insufficient moisture, existing pods are shed and new pods are formed. The optimal moisture content of the soil should be 26-36%. Physiologically, soybean belongs to the group of mesophytes, easily overcomes air and soil drought. Soybean water requirements vary by developmental stage. It requires a lot of water for seed germination and grass germination. The highest water consumption of the plant is observed during flowering and seed filling. During this period, the fastest growth of the stem, the highest average daily growth of the leaf area and the rapid formation of pods are observed. If there is a lack of water during this period, the buds, flowers and nodes of the plant will die. Drought during flowering can reduce seed yield by 50% or more. According to the results of many years of breeding, soybean varieties are well adapted to drought conditions, and in drought conditions, it is possible to get a reliable harvest of 0.5-0.9 t/ha in early varieties and 1.0-1.4 t/ha late ripening varieties were harvested. Soybean has biological adaptation properties to drought conditions: reduction of transpiration, closing of stomata, hairiness. It requires a lot of water for seed germination and grass germination. In drought-resistant species, in conditions of low soil moisture, the amount of water retained in their cells increases, cell sap and osmotic pressure, the maximum temperature of protein coagulation is high, and they differ in the characteristic of high synthesis. flowers, pods and seeds are shed. The shedding of vegetative organs increases resistance to drought, protects against water shortages, shadow formation has been observed under the influence of drought. The flowering period is much longer. The number of pods compared to the number of formed flowers was 43-66% according to the varieties. Among the varieties, the number of woven pods was 46-49% compared to the number of formed pods. The number of ripened seeds per plant varied from 68 (Volna variety) to 115 (Rannyaya-10 variety), their weight was 9.5-14.6 grams. (VF Baranov, VMLukomes, 2005). From flowering to pod formation, water consumption increases, the temperature is high. Water consumption is high in the phases of podding and filling, but the increase in filling with organic matter decreases. Drought in the flowering phase reduced the yield by 48-58%, drought in the



flowering period drops the pods and flowers (87-92%), the seed weight decreases. A moderately developed soybean plant consumes 100-150 g of water per day from germination to flowering, and 300-350 g from flowering to full seed formation. It is during this period that soybean varieties are very sensitive to air and soil drought.

The total amount of water used for the production of a crop unit is called the water absorption (water absorption) coefficient. Soybean varieties also differ according to this indicator. The coefficient of water absorption is lower in water-supplied conditions than in drought conditions, especially in the critical phases of flowering and grain filling.

*Resistance to excess moisture* . When soil moisture increases, root aeration is disturbed, transpiration increases, the amount of chlorophyll in leaves decreases, the productivity of photosynthesis decreases, and grains are not filled. The amount of nitrogen in the plant varies by phase. During the weeding period, the plant does not grow well, aerial roots develop. If these conditions last up to 10 days, the plant will die. After two weeks, when the optimal humidity is restored, the plant continues to develop, but does not completely recover.

### References:

1. Sullieva S., Zokirov K. Biology of weeds //Scientific Bulletin of Namangan State University. – 2019. – Т. 1. – №. 2. – С. 120-124.
2. Фозилов Ш. М. Периодичность роста и формирования урожая у внутривидовых форм пшеницы //Интернаука. – 2019. – №. 45-1. – С. 18-20.
3. Суллиева С. Х., Зокиров К. Г. Вредные свойства и классификация сорняков //Экономика и социум. – 2020. – №. 9 (76). – С. 319-322.
4. Vaxriddinova R. U., Musurmonovich F. S. Soybean-as a source of valuable food //Texas Journal of Multidisciplinary Studies. – 2022. – Т. 6. – С. 165-166.
5. Xurramovna S. S. et al. Use Atlantis Herbicide for Avena Fatua in the Middle of a Wheat Field //International Journal on Integrated Education. – 2023. – Т. 3. – №. 1. – С. 45-47.



6. Normuminovna Q. D., Musurmonovich F. S. Bioecological Properties of *Salvia Officinalis* L //Texas Journal of Multidisciplinary Studies. – 2022. – Т. 6. – С. 249-252.
7. Musurmonovich F. S., Komiljonovna X. S., Qudrat o'g'li S. A. Some Photosynthetic Indicators of Soybean Varieties //Texas Journal of Multidisciplinary Studies. – 2022. – Т. 5. – С. 255-257.
8. Суллиева С. Х., Зокиров К. Г. Структура урожайности озимой пшеницы при применении гербицидов против сорняков //Экономика и социум. – 2020. – №. 9 (76). – С. 323-326.
9. Sullieva S. X., G'Zokirov Q. The Structure of the Yield of Winter Wheat When Using Herbicides Against Weeds //International Journal on Integrated Education. – 2023. – Т. 3. – №. 11. – С. 37-40.
10. Суллиева С. Х., Бобоева Н. Т., Зокиров К. Г. Виды и сорта хризантем //Экономика и социум. – 2019. – №. 10 (65). – С. 315-317.
11. Суллиева С. Х., Бобоева Н. Т., Зокиров К. Г. Эффективность земли вместе в текущем лечении //Мировая наука. – 2019. – №. 6 (27). – С. 450-452.
12. Sullieva S. et al. Effect of planting time and seedling feeding area on the yield and biochemical composition of leek (*Allium porrum* L.) //BIO Web of Conferences. – EDP Sciences, 2024. – Т. 93. – С. 02006.
13. Sulliyeva S., Zokirov Q. Reproduction plantations of (hippophae) in surhandarya region //Scientific Bulletin of Namangan State University. – 2019. – Т. 1. – №. 2. – С. 148-150.
14. Khurramovna S. et al. Technology of growing zaffaron in the surhandarya area //International journal of discourse on innovation, integration and education. – 2020. – Т. 1. – №. 5. – С. 335-337.
15. Суллиева С. Х. и др. Методология планирования классической работы //Экономика и социум. – 2020. – №. 1 (68). – С. 774-776.
16. Суллиева С. Х. Влияние применения гербицидов против сорных растений на урожайность озимой пшеницы в староорошаемых землях сурхандарьинской



- области //Актуальные научные исследования в современном мире. – 2016. – №. 11-2. – С. 26-29.
17. Abduvasikov A. et al. The concept of production resources in agricultural sector and their classification in the case of Uzbekistan //Caspian Journal of Environmental Sciences. – 2024. – Т. 22. – №. 2. – С. 477-488.
18. Khurramovna S. S. et al. Economic efficiency of application of herbicides against double and double weeds in autumn wheat field //European Scholar Journal. – 2021. – Т. 2. – №. 7. – С. 74-76.
19. Sulliyeva S. X. et al. BIOLOGIK TA’LIMNING ROLI //Журнал естественных наук. – 2021. – Т. 1. – №. 1.
20. Суллиева С. Бегона ўтларга қарши кураш самарадорлиги //Ж. Ўзбекистон қишлоқ хўжалиги. – 2008. – №. 5.
21. Khalikov B. et al. Effect of planting time and seedling feeding area on leek (*Allium porrum*) plant biometric parameters //BIO Web of Conferences. – EDP Sciences, 2024. – Т. 93. – С. 02020.
22. Zokirov K., Sullieva S. Effect of storage conditions and periods on the quality of watermelon (*Citrullus lanatus*) //E3S Web of Conferences. – EDP Sciences, 2023. – Т. 452. – С. 01016.
23. Ergashovich K. A., Musurmonovich F. S. Some Characteristics Of Transpiration Of Promising Soybean’s Varieties //The American Journal of Agriculture and Biomedical Engineering. – 2021. – Т. 3. – №. 05. – С. 28-35.
24. Baxriddinovna R. U., Musurmonovich F. S. Distance Learning System in Educational System Instead, and Significance //Texas Journal of Multidisciplinary Studies. – 2023. – Т. 21. – С. 11-13.
25. Fozilov S. Effect of stress factors on some physiological parameters of soybean plant //Science and innovation in the education system. – 2023. – Т. 2. – №. 7. – С. 722-74.
26. Baxriddinovna R. U. Methodology For Solving Problems of Food Chains and Ecological Pyramids and Its Significance //Texas Journal of Multidisciplinary Studies. – 2024. – Т. 28. – С. 19-22.



27. Fozilov S. The effect of drought on the water regime in the leaves of soybean varieties //Science and innovation in the education system. – 2023. – T. 2. – №. 9. – C. 25-28.
28. Ravshanova U. B. et al. Soya o'zbekistonda oziq-ovqat xavfsizligini ta'minlashning muhim manbai sifatida //Экономика и социум. – 2022. – №. 11-1 (102). – C. 81-84.
29. Musurmonovich F. S. et al. Peculiarities of the intensity of photosynthesis and transpiration of soy leaves //Ann. For. Res. – 2022. – T. 65. – №. 1. – C. 5371-5378.
30. Fozilov S., Ziyodova M. Maktablarda steam texnologiyasini joriy etishning xususiyatlari va afzalliklari //Biologiyaning zamonaviy tendensiyalari: muammolar va yechimlar. – 2023. – T. 1. – №. 5. – C. 819-821.
31. Baxriddinova R. U., Musurmonovich F. S. Maktabda tabiiy fanlar, texnologiya, muhandislik, san'at va matematika fanlarini uyg'unlikda o'qitishning afzalliklari //Nazariy va amaliy fanlardagi ustuvor islohotlar va zamonaviy ta'limning innovatsion yo'nalishlari. – 2024. – T. 1. – №. 4. – C. 259-263.
32. Xayitovna P. M., Faxriddinovich M. S. Types of corn grown in Uzbekistan and their peculiarities //Texas Journal of Agriculture and Biological Sciences. – 2022. – T. 3. – C. 59-63.
33. Musurmonovich F. S., Baxriddinova R. U. Soya barglarida fotosintez va transpiratsiya jadalligining o'ziga xos xususiyatlari // Nazariy va amaliy fanlardagi ustuvor islohotlar va zamonaviy ta'limning innovatsion yo'nalishlari. – 2024. – T. 1. – №. 4. – C. 268-272.
34. Musurmonovich F. S., Baxriddinova R. U. Oqsil taqchilligini ta'minlashda soya o'simligining o'rni //nazariy va amaliy fanlardagi ustuvor islohotlar va zamonaviy ta'limning innovatsion yo'nalishlari. – 2024. – T. 1. – №. 4. – C. 254-258.
35. Baxriddinova R. U., Musurmonovich F. S. Oziq zanjiri va oziq to'ri tuzish // Nazariy va amaliy fanlardagi ustuvor islohotlar va zamonaviy ta'limning innovatsion yo'nalishlari. – 2024. – T. 1. – №. 4. – C. 264-267.
36. Musurmonovich F. S. Bug'doy navlarining gullash fazasida suv almashinuv xususiyatlari //Nazariy va amaliy fanlardagi ustuvor islohotlar va zamonaviy ta'limning innovatsion yo'nalishlari. – 2024. – T. 1. – №. 5. – C. 578-583.



37. Eshonqulov S. S. et al. The effectiveness of explained the content of the subject of hybridization to students in school lessons //Экономика и социум. – 2024. – №. 3-1 (118). – С. 163-168.
38. Musurmonovich F. S. Dorivor mavrakning bioekologiyasining o'ziga xos xususiyatlari //Nazariy va amaliy fanlardagi ustuvor islohotlar va zamonaviy ta'limning innovatsion yo'nalishlari. – 2024. – T. 1. – №. 5. – С. 571-577.